



Aura OMI observations of global SO₂ and NO₂ pollution from 2005 to 2013



Nickolay Krotkov¹, Can Li ^{1,2}, Lok Lamsal^{1,3}, Edward Celarier^{1,3},
Sergey Marchenko⁴, William H. Swartz⁵, Eric Bucsela⁶, Vitali
Fioletov⁷, Chris McLinden⁷, Joanna Joiner¹, Pawan K. Bhartia¹, Bryan
Duncan¹, Russ Dickerson⁸

¹NASA Goddard Space Flight Center, ² ESSIC, University of Maryland College Park,
³GESTAR, University Space Research Association, ⁴ Science Systems and
Applications, Inc., ⁵ Applied Physics Laboratory, John Hopkins University, ⁶ SRI
International, ⁷Environment Canada, Ontario CA, ⁸ Department of Atmospheric
and Oceanic Science, University of Maryland College Park

Aura STM College Park, MD

18 September 2014




Key improvements in OMI NO₂ and SO₂



- **Significant improvements in retrieval quality –**
 - Improved spectral fitting of OMI NO₂ removes 20%- 40% of the stratospheric biases with other satellite measurements. New NO₂ version planned for release next year
 - New PCA SO₂ algorithm uses full spectral content from OMI, reduces noise by half and removes biases (artifacts)
 - New Version 2 OMI SO₂ dataset will be released this fall
- **Maximal data continuity between instruments –**
 - Both OMI NO₂ and SO₂ algorithms can benefit new missions: SNPP/ OMPS, TROPOMI, GEMS and TEMPO
 - no need to develop instrument-specific radiance data correction schemes
- **Maximal sensitivity -**
 - PCA SO₂ detection limit for point sources is half the current PBL algorithm
- **Flexibility –**
 - PCA SO₂ fitting window can be easily adjusted to optimize sensitivity under different conditions: from small anthropogenic signals to largest volcanic plumes.
 - NO₂ fitting window can be expanded to UV wavelengths (OMPS)

Regional trends in OMI new SO₂ and NO₂ : 2005-2013



OMI SO₂ and NO₂ time series

- SZA < 70°
 - Cross-track CCD rows 6-23 (excluding row anomaly for all years) ;
 - Snow-free observations (according to the IMS data* product);
 - SCD-O₃<1500 DU, VCD_SO₂<15 DU
 - Additional volcanic filtering: all days removed which, over that region and considering all years, had a daily 99.9th percentile value greater than X,
 - where X=5 DU for Eastern North America,
8 DU for Eastern Europe and India,
10 DU for China –these thresholds are obtained using the 99.9 percentile daily regional time series.
- For consistency removed the same volcanic days in NO₂ product



* Interactive multi-sensor snow and ice product, <http://www.natice.noaa.gov/ims/>

2005-2007



Eastern US

SO₂

2011-2013





NO₂





SO₂

Eastern US



NO₂





Eastern Europe

2005-2007

2011-2013

SO₂





Etna Volcano

Eastern Europe



2005-2007

2011-2013


NO₂





Eastern Europe: Time series for Maritsa Iztok



2005



2013





SO₂

2005-2007




India

2011-2013



Chhattisgarh

Pronunciation: chuht-tihs-guhr




NO₂


2005-2007

India

2011-2013






Power plants / smelter





Chhattisgarh

Pronunciation: chuht-tihs-guhr





Time series: India (Chhattisgarh)

SO₂




NO₂




Eastern Asia

2005-2007



2011-2013





Volcano

Eastern Asia



2005-2007

2011-2013





Eastern China: Time series

SO₂




NO₂




Summary


Eastern USA



Eastern Europe (Maritsa Iztok, Bulgaria)



India (Chhattisgarh)



Eastern Asia (Eastern China)

